

CHM 305 - Lecture 5 - The Rules of Quantum Mechanics (Part 1)

Prof. Marissa Weichman

We are now going to work towards the big rules of the game, which are usually referred to as the “Postulates of Quantum Mechanics.” We will start to get there this class and by the end of next class. This material is covered in Chapter 4 of McQuarrie.

1 Quantum Operators

In quantum mechanics, every measurable physical quantity (e.g. energy, momentum, position) is called an *observable*. Each observable has a corresponding operator, which we denote with a caret like \hat{O} . Operators simply tell you to “do something” to whatever follows them. We already know some examples of operators:

- $\hat{H} = \frac{-\hbar^2}{2m} \frac{\partial^2}{\partial x^2} + V(x)$, the Hamiltonian, or energy operator, in one dimension
- $\hat{x} = x$, the position operator is simply multiplication by x
- $\hat{p} = -i\hbar \frac{\partial}{\partial x}$, the momentum operator, closely related to differentiation by x . (See the note at the end of the lecture 4 notes for more on where this definition of \hat{p} comes from.)

1.1 Eigenfunctions and Eigenvalues

When we solve the time-independent Schrödinger equation

$$\hat{H}\psi_n = E_n\psi_n \tag{1}$$

we are actually solving an *eigenvalue problem* where the eigenfunctions ψ_n are a set of special functions that when acted on by that operator return themselves times a number. And the special number E_n is the eigenvalue that corresponds to the n^{th} eigenfunction. If you are familiar with linear algebra, a useful analogy is that operators can be represented by matrices with eigenfunctions represented by vectors.

Note: There is an entire formalism of quantum mechanics based around matrices, which we will only touch on in this course. If you take graduate level quantum mechanics you will see a lot more of that.

In quantum mechanics, eigenvalues are always real and represent the measured value of the observable. Using the Hamiltonian operator as an example, our observable is energy, and the eigenvalues E_n represent the energy of the system when its state is a given stationary state (or eigenstate) ψ_n .

Practice Problem 1: Check if $\psi(x) = \sin(kx)$ is an eigenfunction of the following operators:

$$\hat{A} = \frac{d}{dx} \quad (2)$$

$$\hat{B} = \frac{d^2}{dx^2} \quad (3)$$

If you find $\psi(x)$ to be an eigenvector of one of these operators, what is its eigenvalue?

1.2 Quantum operators are linear

If \hat{A} is an operator with eigenfunctions ψ_1 and ψ_2 and corresponding eigenvalues a_1 and a_2 , then:

$$\hat{A}(c_1\psi_1 + c_2\psi_2) = c_1a_1\psi_1 + c_2a_2\psi_2 \quad (4)$$

assuming c_1 and c_2 are constants. In other words, \hat{A} acts independently on each function in turn.

To further generalize this phenomenon, if $\{\psi_n\}$ are the set of eigenfunctions of \hat{A} with corresponding eigenvalues $\{a_n\}$ then:

$$\hat{A} \sum_{i=1}^{\infty} c_i \psi_i = \sum_{i=1}^{\infty} c_i a_i \psi_i \quad (5)$$

1.3 Quantum operators need not commute

As with matrix multiplication, it's important to act with operators in the correct order. Whichever operator is nearest to the wavefunction always acts first:

$$\hat{A}\hat{B}\psi(x) = \hat{A}\left(\hat{B}\psi(x)\right) \quad (6)$$

Another way to say this is that operators do not necessarily commute, so in general:

$$\hat{A}\hat{B}\psi(x) \neq \hat{B}\hat{A}\psi(x) \quad (7)$$

In fact, in the rare cases where two operators *do* commute with one another, it implies that they have a very special relationship, which we will learn more about later.

We will also use something called a *commutator* as shorthand:

$$[\hat{A}, \hat{B}] \psi \equiv (\hat{A}\hat{B} - \hat{B}\hat{A})\psi \quad (8)$$

where $[\hat{A}, \hat{B}] = 0$ if the operators commute and $[\hat{A}, \hat{B}] \neq 0$ if they do not.

Practice Problem 2: Do the operators $\hat{A} = \frac{d}{dx}$ and $\hat{B} = x$ commute? *Hint: evaluate $[\hat{A}, \hat{B}]\psi$.*

2 Eigenfunctions of Quantum Operators

Let's hammer out a few rules for how eigenfunctions of quantum mechanical operators should behave. Eigenfunctions have some peculiar properties that we'll need to make use of as we proceed.

Consider again the set of eigenfunctions of the operator \hat{A} , which we will call $\{\psi_n(x)\}$. By definition, the eigenfunctions have the property that when operated on by \hat{A} , they return themselves times a constant, their eigenvalue:

$$\hat{A}\psi_n(x) = a_n\psi_n(x) \quad (9)$$

2.1 Eigenfunctions are Normalized

For each eigenfunction $\psi_n(x)$ we will mandate that the integral of the eigenfunction's magnitude over all space be normalized to 1:

$$\int_{-\infty}^{\infty} \psi_n^*(x)\psi_n(x)dx = \int_{-\infty}^{\infty} |\psi_n(x)|^2 dx = 1 \quad (10)$$

Note: $\psi_n^*(x)$ is the complex conjugate of $\psi_n(x)$. If $\psi_n(x)$ is real, then $\psi_n^*(x) = \psi_n(x)$. Why do we need to take the complex conjugate? $\psi_n(x)$ may be complex, and the magnitude of a complex number $y = a + ib$ is given by $|y|^2 = y^* \cdot y = (a - ib) \cdot (a + ib)$.

Normalization is important because wavefunctions represent probability distributions for where we expect to find a particle. We want the total probability that the particle be found *somewhere* in space to be 1.

Practice Problem 3: Normalize the wavefunction $\psi'(x) = 1 - x$ over the interval $[0, 1]$.

2.2 Eigenfunctions of the Same Operator are Orthogonal


Consider again our infinite set of functions $\psi_n(x)$, which are all eigenfunctions of the same operator \hat{A} . It turns out that any pair of eigenfunctions with $m \neq n$ are orthogonal. Two functions are by definition orthogonal if:

$$\int_{-\infty}^{\infty} \psi_n^*(x)\psi_m(x)dx = 0 \quad (11)$$

What this means physically is that each eigenfunction must be independent of the others - they do not interfere or overlap. As an example, the set of functions


$$\psi_n(x) = \sin(2\pi nx) \quad n = 1, 2, 3, \dots \quad (12)$$

are all mutually orthogonal. Let's plot the first two of these functions over the interval $x = [0, 1]$:



You should be able to convince yourself that the net overlap of these two functions is exactly 0 over the interval shown. In particular, if you consider the product of these two functions as plotted in the righthand panel above, by inspection we can see that the integral of this quantity will vanish over one complete oscillation.

Another good example of orthogonal functions are atomic orbitals. Visualize an atomic s orbital and a p orbital. Because the p orbital has two nodes with amplitudes of opposite sign, its net overlap with the s orbital over all space is 0. Any pair of atomic orbitals you might choose are similarly orthogonal over all (3D) space.



2.3 Eigenfunctions of the Same Operator Form a Complete Set

We will often construct wavefunctions out of linear combinations of “basis functions.” To give a familiar example, we build molecular orbitals as linear combinations of atomic orbitals.

If our basis functions $\{\psi_n(x)\}$ are the infinite ladder of eigenfunctions for a quantum mechanical operator \hat{A} , then they form a *complete basis set*. This means that for *any* arbitrary wavefunction $\Phi(x)$, we can find coefficients b_n such that:

$$\Phi(x) = \sum_{n=1}^{\infty} b_n \psi_n(x) \tag{13}$$

How do we go about finding the correct set of coefficients b_n to represent the desired $\Phi(x)$? We can actually derive an expression very simply, using the orthogonality relation we just discussed for eigenfunctions. Consider the following integral, which we can think of as an *overlap* integral between $\Phi(x)$ and the m^{th} eigenfunction:

$$\int_{-\infty}^{+\infty} \psi_m^*(x) \Phi(x) dx \quad (14)$$

$$= \int_{-\infty}^{+\infty} \psi_m^*(x) \left(\sum_{n=1}^{\infty} b_n \psi_n(x) \right) dx \quad (15)$$

$$= \sum_{n=1}^{\infty} b_n \int_{-\infty}^{+\infty} \psi_m^*(x) \psi_n(x) dx \quad (16)$$

$$= \sum_{n=1}^{\infty} b_n \delta_{mn} = b_m \quad \checkmark \quad (17)$$

where

$$\delta_{mn} = \begin{cases} 0, & m \neq n \\ 1, & m = n \end{cases} \quad (18)$$

is a useful shorthand called the Kronecker delta.

So it proves to be quite straightforward to find the set of b_n coefficients and express $\Phi(x)$ as an expansion of eigenfunctions. This is a very important and useful property of eigenfunctions. This is a concept closely related to Fourier series, if you have seen that in previous coursework.

Practice Problem 4: (a) Normalize the set of functions $\psi_n(x) = Ne^{inx}$, over the window $0 \leq x \leq 2\pi$. To do so, you will need to find the normalization constant N such that:

$$\int_0^{2\pi} \psi_n^*(x) \psi_n(x) dx = 1$$

where $\psi_n^*(x)$ is the complex conjugate of $\psi_n(x)$.

(b) Show that your normalized set of functions $\psi_n(x) = Ne^{inx}$ is orthonormal if n is an integer. To do so, you will need to show that the integral

$$\int_0^{2\pi} \psi_m^*(x) \psi_n(x) dx = \begin{cases} 1 & \text{if } m = n \\ 0 & \text{if } m \neq n \end{cases}$$