CHM 502 – Module 2 – An Aside on Time Evolution

Prof. Marissa Weichman

Let's take a moment to talk about time-dependent quantum mechanics in a more general way, and introduce some machinery that will be useful to us soon. Much of this material was inspired by notes from Andrei Tokmakoff at UChicago.

1 The time evolution operator (for $\hat{H} \neq \hat{H}(t)$)

We've talked about how the wavefunction of a system evolves according to the time-dependent Schrödinger equation (TDSE):

$$i\hbar \frac{d}{dt} |\Psi(t)\rangle = \hat{H} |\Psi(t)\rangle \tag{1}$$

and how we can expand a general time-dependent wavefunction in the basis of energy eigenstates:

$$|\Psi(t)\rangle = \sum_n c_n(0)\ e^{-iE_n t/\hbar}\ |\psi_n\rangle \tag{2}$$

Assuming for the moment that we have a time-independent Hamiltonian, we can manipulate (or "integrate") the TDSE to find:

$$\frac{d}{dt} |\Psi(t)\rangle = -\frac{i\hat{H}}{\hbar} |\Psi(t)\rangle \tag{3}$$

$$\rightarrow \quad |\Psi(t)\rangle = e^{-i\hat{H}t/\hbar}\ |\Psi(0)\rangle \equiv \hat{U}(t,0)\ |\Psi(0)\rangle \tag{4}$$

We are arguing here that if the temporal derivative of $|\Psi(t)\rangle$ returns a quantity proportional to itself, then we can quickly write down a solution to the differential equation, making use of the properties of complex exponentials.

In the meantime, we have defined $\hat{U}(t,0)$ as the *time evolution* operator or *propagator*, whose application evolves our wavefunction forward in time from $|\Psi(0)\rangle$ to $|\Psi(t)\rangle$.

What does it mean to have the $\hat{H}$ operator (or any operator for that matter) living inside a complex exponential? We can write this down explicitly as a Taylor expansion in powers of that operator. Let's check that our expression for $|\Psi(t)\rangle$ in Eqn. 4 behaves as we expect:

$$\hat{U}(t,0)\,|\Psi(t)\rangle = e^{-i\hat{H}t/\hbar}\,|\Psi(0)\rangle \tag{5}$$

$$= \left[1 - \frac{i\hat{H}t}{\hbar} + \frac{1}{2}\left(\frac{-i\hat{H}t}{\hbar}\right)^2 + \cdots\right] \sum_n c_n(0)\,|\psi_n\rangle \tag{6}$$

$$= \sum_n c_n(0) \left[1 - \frac{i\hat{H}t}{\hbar} + \frac{1}{2}\left(\frac{-i\hat{H}t}{\hbar}\right)^2 + \cdots\right] |\psi_n\rangle \tag{7}$$

$$= \sum_n c_n(0) \left[1 - \frac{iE_nt}{\hbar} + \frac{1}{2}\left(\frac{-iE_nt}{\hbar}\right)^2 + \cdots\right] |\psi_n\rangle \tag{8}$$

$$= \sum_n c_n(0)\, e^{-iE_nt/\hbar}\,|\psi_n\rangle = |\Psi(t)\rangle \qquad \checkmark \tag{9}$$

A few more notes on how this $\hat{U}(t,0)$ beast behaves:

- $\hat{U}(t,0)$ acts to the right on *kets* to evolve forward in time
- $\hat{U}^\dagger(t,0)$ acts to the left on *bras* to evolve forward in time. However, one can easily show that

  $$\hat{U}^\dagger(t,0) = e^{i\hat{H}t/\hbar} = \hat{U}^{-1}(t,0) \tag{10}$$

  so the $\hat{U}^\dagger(t,0)$ operator will also act to the right on on *kets* to evolve them backwards in time. $\hat{U}^\dagger(t,0)$ is therefore also referred to as the time reversal operator.
- $\hat{U}(t,0)$ is unitary, meaning that $\hat{U}^\dagger(t,0)\,\hat{U}(t,0) = \hat{I}$. This ensures that the normalization of a wavefunction is preserved as it propagates in time.
- Finally, the time evolution operator has the nice property that:

  $$\hat{U}(t_2,t_0) = \hat{U}(t_2,t_1)\,\hat{U}(t_1,t_0) \tag{11}$$

  Note that order of operation is very important here. In general, $\hat{U}$ does not commute with itself at different times:

  $$\hat{U}(t_2,t_1)\,\hat{U}(t_1,t_0) \neq \hat{U}(t_1,t_0)\,\hat{U}(t_2,t_1) \tag{12}$$

2 The time evolution operator for $\hat{H} = \hat{H}(t)$

We assumed earlier that our Hamiltonian is time-independent, particularly in going from Eqn. 3 to Eqn. 4. What does the time evolution operator look like for a *time-dependent* Hamiltonian? Let's go back to the TDSE to investigate.

$$i\hbar\frac{d}{dt}\left|\Psi(t)\right\rangle = \hat{H}(t)\left|\Psi(t)\right\rangle \tag{13}$$

$$i\hbar\frac{d}{dt}\left[\hat{U}(t,0)\left|\Psi(0)\right\rangle\right] = \hat{H}(t)\left[\hat{U}(t,0)\left|\Psi(0)\right\rangle\right] \tag{14}$$

$$\left[i\hbar\frac{d}{dt}\hat{U}(t,0)\right]\left|\Psi(0)\right\rangle = \left[\hat{H}(t)\,\hat{U}(t,0)\right]\left|\Psi(0)\right\rangle \tag{15}$$

The two operators acting on $|\Psi(0)\rangle$ on either side of the equals sign above must be equivalent, so:

$$i\hbar\frac{d}{dt}\hat{U}(t,0) = \hat{H}(t)\,\hat{U}(t,0) \tag{16}$$

***Aside*:** Why not just directly integrate Eqn. 16? Let's see what happens and why it turns out to be problematic:

$$i\hbar\frac{d}{dt}\hat{U}(t,0) = \hat{H}(t)\hat{U}(t,0) \tag{17}$$

$$\frac{d\hat{U}(t,0)}{U(t,0)} = -\frac{i}{\hbar}\hat{H}(t)\,dt \tag{18}$$

$$\int_{U(t_0,0)}^{U(t,0)}\frac{d\hat{U}(t',0)}{U(t',0)} = \ln\left[U(t,0)\right] - \ln\left[U(t_0,0)\right] = -\frac{i}{\hbar}\int_{t_0}^{t}dt'\hat{H}(t') \tag{19}$$

$$\frac{U(t,0)}{U(t_0,0)} = \frac{U(t,t_0)\cancel{U(t_0,0)}}{\cancel{U(t_0,0)}} = U(t,t_0) = \exp\left[-\frac{i}{\hbar}\int_{t_0}^{t}dt'\hat{H}(t')\right] \tag{20}$$

We can now attempt to evaluate the RHS complex exponential using a Taylor expansion:

$$U(t,t_0) \stackrel{?}{=} 1 - \frac{i}{\hbar}\int_{t_0}^{t}dt'\,\hat{H}(t') + \frac{1}{2!}\left(-\frac{i}{\hbar}\right)^2\left[\int_{t_0}^{t}dt'\,\hat{H}(t')\right]^2 + \cdots \tag{21}$$

Let's inspect the second term of this expansion a little more closely:

$$\left[\int_{t_0}^{t}dt'\,\hat{H}(t')\right]^2 = \left[\int_{t_0}^{t}dt'\,\hat{H}(t')\right]\cdot\left[\int_{t_0}^{t}dt''\,\hat{H}(t'')\right] = \int_{t_0}^{t}\int_{t_0}^{t}dt'dt''\,\hat{H}(t')\hat{H}(t'') \tag{22}$$

And now we see the problem: this expression integrates factors of $\hat{H}(t')$ and $\hat{H}(t'')$ over a range of times that scrambles their relative time orderings. Unfortunately, we are not guaranteed that the Hamiltonian commutes with itself at different times: $[\hat{H}(t'), \hat{H}(t'')] \neq 0$. We have not treated $\hat{H}$ properly as an operator, so none of the math in this aside is valid!

Let's proceed a bit more carefully instead and see what happens if we integrate Eqn. 16 without rearranging it. We will integrate both sides with respect to time using τ as a dummy variable for integration:

$$i\hbar \int_0^t \frac{d}{d\tau}\hat{U}(\tau,0)d\tau = \int_0^t \hat{H}(\tau)\hat{U}(\tau,0)d\tau \tag{23}$$

$$= i\hbar \left[\hat{U}(\tau,0)\right]_0^t = i\hbar \left[\hat{U}(t,0) - \hat{U}(0,0)\right] = i\hbar \left[\hat{U}(t,0) - 1\right] \tag{24}$$

$$\rightarrow \quad \boxed{\hat{U}(t,0) = 1 - \frac{i}{\hbar}\int_0^t d\tau\, \hat{H}(\tau)\,\hat{U}(\tau,0)} \tag{25}$$

This is a bit of a weird expression, since it casts $\hat{U}(t,0)$ in terms of an integral involving itself. But we can work with it by iteratively substituting $\hat{U}(t,0)$ into the above, e.g.

$$\hat{U}(t,0) = 1 - \frac{i}{\hbar}\int_0^t d\tau\, \hat{H}(\tau)\left[1 - \frac{i}{\hbar}\int_0^\tau d\tau'\, \hat{H}(\tau')\hat{U}(\tau',0)\right] \tag{26}$$

$$= 1 \;+\; \left(\frac{-i}{\hbar}\right)\int_0^t d\tau\, \hat{H}(\tau) \;+\; \left(\frac{-i}{\hbar}\right)^2 \int_0^t d\tau \int_0^\tau d\tau'\, \hat{H}(\tau)\,\hat{H}(\tau')\,\hat{U}(\tau',0) \tag{27}$$

$$= \ldots$$

$$= 1 \;+\; \sum_{n=1}^{\infty}\left(\frac{-i}{\hbar}\right)^n \int_0^t d\tau_n \int_0^{\tau_n} d\tau_{n-1} \ldots \int_0^{\tau_2} d\tau_1\, \hat{H}(\tau_n)\hat{H}(\tau_{n-1}) \;\ldots\; \hat{H}(\tau_2)\,\hat{H}(\tau_1) \tag{28}$$

Note that there is a time-ordering to each of these terms such that $t \geq \tau \geq \tau' \geq \tau'' \ldots$ and $t \geq \tau_n \geq \cdots \geq \tau_2 \geq \tau_1$. By keeping careful track of this time ordering, we avoid the pitfalls discussed in the aside above.

You can perform a sanity check for yourself that taking $\hat{H} \neq \hat{H}(t)$, Eqn. 28 gives us back Eqn. 21, as a time-independent Hamiltonian *does* commute with itself. In addition, evaluating Eqn. 21 for $\hat{H} \neq \hat{H}(t)$ also gives us back our definition of $\hat{U}(t,0)$ from Eqn. 4.

It can be difficult to conceptualize what is going on here. One way to think about it is that these nested integrals are counting up all the possible ways the time evaluation operator can take us from initial state $|\Psi(0)\rangle$ to final state $|\Psi(t)\rangle$, as both our wavefunction and the Hamiltonian change in time. The different trajectories, represented by different terms in the sum, are mediated by visits to all possible intermediate states of the system at all possible intermediate times. All of these different paths interfere with each other as they acquire their own phases.

That's all we will say about this machinery for the time being. It crops up when you are considering the dynamics of systems that are perturbed with a time-dependent Hamiltonian at various points in time, as you might have in a nonlinear spectroscopy experiment where time-ordered pulses interact with the system. More on this later.

3 The "pictures" of time-dependent quantum mechanics

Now that we have introduced the time evolution operator, it's a good time to make one more aside about the various formalisms for thinking about time-dependent quantum mechanics.

3.1 The Schrödinger picture

So far, we have worked within the "Schrödinger picture," where wavefunctions are time-dependent, and are propagated in time as:

$$|\Psi(t)\rangle = \hat{U}(t,0)\,|\Psi(0)\rangle \tag{29}$$

Meanwhile, we tend to think of operators as time-independent quantities:

$$\hat{x} = x \neq \hat{x}(t) \tag{30}$$

$$\hat{p} = -i\hbar\frac{d}{dx} \neq \hat{p}(t) \tag{31}$$

While the expectation values of these operators may evolve with time, we ascribe this to the time-dependent behavior of the wavefunction:

$$\langle\hat{A}(t)\rangle = \langle\Psi(t)|\hat{A}|\Psi(t)\rangle \tag{32}$$

$$\equiv \langle\Psi_S(t)|\hat{A}_S|\Psi_S(t)\rangle \tag{33}$$

where the subscript "S" denotes wavefunctions and operators defined in the Schrödinger picture.

3.2 The Heisenberg picture

Let's now introduce the "Heisenberg picture," in which operators are time-propagated while wavefunctions are time-independent. This turns out to be entirely equivalent to the Schrödinger picture.

Consider:

$$\langle\hat{A}(t)\rangle = \langle\Psi_S(t)|\hat{A}_S|\Psi_S(t)\rangle \tag{34}$$

$$= \langle\Psi_S(0)|\hat{U}^\dagger(t,0)\hat{A}_S\hat{U}(t,0)|\Psi_S(0)\rangle \tag{35}$$

$$\equiv \langle\Psi_H|\hat{A}_H(t)|\Psi_H\rangle \tag{36}$$

where we've used "H" subscripts to denote the Heisenberg picture:

$$\hat{A}_H(t) \equiv \hat{U}^\dagger(t,0)\,\hat{A}_S\,\hat{U}(t,0); \qquad \hat{A}_S = \hat{A}_H(0) \tag{37}$$

$$|\Psi_H\rangle \equiv |\Psi_S(0)\rangle\,; \qquad |\Psi_S(t)\rangle = \hat{U}(t,0)\,|\Psi_H\rangle \tag{38}$$

The Heisenberg wavefunction is for all times the initial, stationary Schrödinger wavefunction at time zero. Meanwhile, the physical observables of the particle (represented by operators) evolve in time – this admittedly makes some physical sense! The Heisenberg picture is sometimes more convenient to work with mathematically than the Schrödinger picture; we'll see it crop up periodically.

3.3 The interaction picture

There is a third useful representation called the "interaction picture" which is often invoked for problems with time-dependent Hamiltonians that can be written as a time-independent reference Hamiltonian plus a small time-dependent perturbation:

$$\hat{H}(t) = \hat{H}_0 + \hat{V}(t) \tag{39}$$

where $\hat{H}_0$ is a relatively simple system that we can solve exactly, while $\hat{V}(t)$ may be quite complicated.

We can define a time-evolution operator that involves just the reference Hamiltonian:

$$\hat{U}_0(t,0) = e^{-i\hat{H}_0 t/\hbar} \tag{40}$$

We relate the interaction picture to the Schrödinger as follows:

$$|\psi_S(t)\rangle = \hat{U}_0(t,0)\,|\psi_I(t)\rangle \quad \rightarrow \quad \boxed{|\psi_I(t)\rangle = \hat{U}_0^{-1}(t,0)\,|\psi_S(t)\rangle} \tag{41}$$

The interaction representation defines wavefunctions in such a way that we strip away the (uninteresting) $e^{-i\hat{H}_0 t/\hbar}$ modulation that arises from the time-independent part of the wavefunction, $\hat{H}_0$. This leaves behind a representation of the wavefunction $|\psi_I(t)\rangle$ whose only-time dependence arises from the (much more interesting) time-dependence of the perturbation $V(t)$. More on this later!